

N-049180-S-0021  
 Amprolium and Ethopabate/Bacitracin Methylene Disalicylate 5-35/Roxarsone  
 Broiler Chicken Ration - WG/COCCI  
 Type C Med. Feed

### Type C Medicated Feed

For increased rate of weight gain and as an aid in the prevention of coccidiosis where severe exposure to coccidiosis from *Eimeria acervulina*, *E. maxima* and *E. brunetti* is likely to occur in broiler chickens raised in floor pens.

### ACTIVE DRUG INGREDIENTS

Amprolium .....	113.5 g/ton
Ethopabate .....	36.3 g/ton
Bacitracin Methylene Disalicylate.....	5-35 g/ton*
Roxarsone.....	34 g/ton

### GUARANTEED ANALYSIS

Crude Protein, not less than.....	%
Lysine, not less than .....	%
Methionine, not less than.....	%
Crude fat, not less than.....	%
Crude fiber, not more than.....	%
Calcium, not less than.....	%
Calcium, not more than .....	%
Phosphorus, not less than .....	%
Salt <sup>1</sup> , not less than.....	%
Salt <sup>1</sup> , not more than .....	%
Sodium <sup>2</sup> , not less than .....	%
Sodium <sup>2</sup> , not more than .....	%

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials

### DIRECTIONS FOR USE

Feed as the sole ration from time chickens are placed on litter until past the time when coccidiosis is ordinarily a hazard.

**CAUTION:** Do not feed to laying chickens. Do not use as a treatment for outbreaks of coccidiosis.

**WARNING:** Withdraw 5 days before slaughter. Use as the sole source of amprolium and organic arsenic.

\*The final finished feed label should list only a single drug concentration.

Net Weight \_\_\_\_\_ lb. (\_\_\_\_\_ kg) on bag or bulk

Manufactured by  
 BlueBird Feed Mill  
 Anytown, Anystate 12345